


Published by Nigerian Society of Physical Sciences. Hosted by FLAYOO Publishing House LTD

Recent Advances in Natural Sciences

Journal Homepage: <https://flayoophl.com/journals/index.php/rans>

Proximate Analysis of *Telfairia Occidentalis* (Fluted Pumpkin) and *Telfairia Pedata* (Oyster Nut) Leaves Consumed in Katsina Metropolis: A Comparative Study

Yahaya Auwal^{a,1,*}, Magaji Ilu Barde^a, Naziru Imam^b, Abdulhamid Murtala^a^aDepartment of Pure & Industrial Chemistry, Faculty of Natural and Applied Sciences, Umaru Musa Yaradua University, Katsina, Nigeria^bDepartment of Industrial Chemistry, Faculty of Science, Federal University Oye-Ekiti, Ekiti State, Nigeria.

ARTICLE INFO

Article history:

Received: 22 July 2023

Received in revised form: 19 August 2023

Accepted: 20 August 2023

Available online: 25 August 2023

Keywords: *Telfairia occidentalis*, Fluted pumpkin, *Telfairia pedata*, Oyster nut, Proximate analysis, Katsina.

DOI:10.61298/rans.2023.1.1.8

ABSTRACT

Telfairia occidentalis and *Telfairia pedata* are highly valued for their nutritious seeds and are consumed widely in sub-Saharan Africa. However, the potential application of their leaves was not thoroughly investigated and documented. Therefore, proximate analysis of their leaves for potential applications is sacrosanct. Samples of the leaves were obtained from the Katsina metropolis and analysed for their proximate compositions using standard analytical procedures. The result indicates that the percentage carbohydrate, crude fibres, and moisture contents values for *Telfairia pedata* and *Telfairia occidentalis* leaves were 49.23 ± 0.79 , 19.50 ± 0.31 , 62.06 ± 0.49 and 49.44 ± 0.55 , 14.68 ± 1.04 , and 71.40 ± 0.31 , respectively. The crude protein, crude fat, and ash contents were 16.04 ± 0.51 , 2.00 ± 0.50 , 13.33 ± 0.31 and 22.45 ± 0.50 , 4.80 ± 0.34 , and 8.60 ± 0.21 for *Telfairia pedata* and *Telfairia occidentalis*, respectively. The crude fibre and ash content were higher in *Telfairia Pedata* than in *Telfairia Occidentalis* leaves. There was no significant difference in the carbohydrate content of the leaves. However, the crude protein, moisture content, and lipid were higher in the *Telfairia Occidentalis* leaf. The leaves could be a good carbohydrate, protein, and dietary fibre source. However, further study is recommended on the leaves' anti-nutrients, mineral and phytochemical analyses.

© 2023 The Author(s). Production and Hosting by FLAYOO Publishing House LTD on Behalf of the Nigerian Society of Physical Sciences (NSPS). Peer review under the responsibility of NSPS. This is an open access article under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

1. INTRODUCTION

Vegetables refer to the consumable components of herbaceous plants that can be consumed either in their entirety or in portions, raw or cooked. They are commonly incorporated into main courses or salads. These vegetables can possess various flavours, ranging from aromatic and bitter to relatively neutral taste. Sunmonu *et al.* [1] reported that these ingredients are commonly consumed in the dietary habits of the typical Nigerian popula-

tion and offer significant quantities of essential minerals. Although leafy vegetables primarily consist of water, they serve as a rich source of minerals, vitamins, and phytochemicals, resembling a comprehensive natural repository. The *Telfairia* genus, classified under the Cucurbitaceae family, encompasses two distinct species found in Africa: *Telfairia occidentalis* Hooker and *Telfairia pedata*, originating from West Africa and East Africa, respectively [2]. Due to its significant value, *Telfairia Occidentalis*, a herbaceous perennial plant, is cultivated annually in some areas of West Africa [3]. *Telfairia occidentalis* is widely cultivated in the southern region of Nigeria, particularly by the Ig-

*Corresponding author: Tel.: +234-703-1242-037;

e-mail: yahyaaauwal1159@gmail.com (Yahaya Auwal)

bos. This vegetable crop has gained significant importance in its dietary practices and is now a prevalent component in regional households [2]. According to Kayode *et al.* [4], traditional medicine employs this substance to enhance blood circulation, bolster the immune system, and alleviate digestive issues and convulsions. Multiple authors have extensively documented the antioxidant property of *Telfairia Occidentalis* [5, 6]. From a nutritional standpoint, this particular leafy vegetable possesses a significant abundance of minerals, including iron, potassium, sodium, phosphorous, calcium, magnesium, and certain essential amino acids [7].

Additionally, it contains notable phytochemicals such as tannins, saponins, flavonoids, and phenolics [4]. *Telfairia pedata*, commonly known as oyster nut, is a perennial plant cultivated in Central and East Africa. According to Alegbejo [8], the plant in question yields seeds characterised by their significant size, elongated shape, and flat structure. Additionally, when subjected to roasting, these seeds exhibit a flavour reminiscent of almonds. *Telfairia pedata* is a native forest food plant in the Uluguru North and West Usambara Mountain regions, known for providing edible nuts and seeds [9]. The seeds of *Telfairia pedata* are utilised in tropical East Africa for oil production and are subjected to roasting and consumption.

Comparative analysis of the nutritional composition of Cucurbitaceae leaves within a specific region is crucial, as it is a key indicator of potential differences that may arise from climatic factors and soil composition. The nutritional value of *Telfairia occidentalis* leaves has been reported in the literature and is widely consumed throughout sub-Saharan Africa. However, the potential of *Telfairia pedata* leaves has not been comparatively investigated. To fill this knowledge gap, there is a need to evaluate the proximate composition of *Telfairia Occidentalis* and *Telfairia pedata* leaves. This will not only increase the understanding of the nutritional content of these leaves but also contribute to the proximate composition database of Cucurbitaceae leaves.

2. METHODOLOGY

Two different types of freshly harvested *Telfairia Hooker* F. leaves were collected from Katsina City, Nigeria, namely *Telfairia occidentalis* and *Telfairia pedata*. The leaves were carefully placed in a clean polyethylene bag for packaging. The leaf samples underwent authentication and identification at the Herbarium Unit of the Department of Biology, Umaru Musa Yar'adua University, located in Katsina, Nigeria. Before commencing the study, the leaves were subjected to a comprehensive purification procedure, which entailed using water to eradicate any residual soil particles. The leaves were thoroughly washed using distilled water and subsequently allowed to undergo a natural drying process for three days within a sheltered environment. Subsequently, the leaves were ground into a fine powder using a mortar and pestle, and the resulting powder was then filtered through a mesh with a pore size of 20 μm . The acquired powder was afterward placed in a plastic container in anticipation of the proximate analysis. In order to ascertain the moisture content, the investigation utilised leaf samples that were freshly gathered. The proximate analysis was performed in accordance with the protocol specified by the Association of Official Analytical Chemists (AOAC) [10].




Figure 1. Comparative proximate values of *Telfairia pedata* and *Occidentalis* leaves

3. RESULTS AND DISCUSSION

The proximate composition of the two leaves is presented in Table 3. In terms of moisture content, the result shows a high percentage of mean moisture content in both leaves, with *Telfairia pedata* slightly higher than *Telfairia occidentalis*. The relatively higher moisture content in the leaves suggests that the vegetables may not be stored for long due to higher water activity. Higher moisture content helps maintain the cell's protoplasmic content [11]. Additionally, it facilitates enhanced functionality of hydrophilic enzymes and coenzymes essential for the metabolic processes of leafy green vegetables [12]. Nevertheless, the elevated moisture content could give rise to a quality issue, as the samples may exhibit increased vulnerability to bacterial spoilage throughout the storage period [13].

Comparatively, from the present analysis, *Telfairia occidentalis* leaf is more susceptible to deterioration by bacteria on storage and easily digestible on consumption than the *Telfairia Pedata* leaf because it contains higher moisture content (Figure 1). Omimakinde *et al.* [14] reported 8.75% of moisture content in fluted pumpkins, similar to but slightly less than in this study. This may be attributable to the plant's geographical location and the research's analytical accuracy. However, the 3.09 ± 0.93 percentage moisture content of *Telfairia pedata* (Oyster Nut) seeds obtained by Mwakasege *et al.* [15] is about five-fold less than what was obtained in this study. The result shows that the leaves significantly differ in percentage moisture content at ($P < 0.05$).

Similarly, relatively high ash content values of 13.30 ± 0.31 and 8.60 ± 0.20 for *Telfairia pedata* and *Telfairia Occidentalis* leaves, respectively, were recorded. Higher ash content is an indication of abundant mineral elements in the leaves. Comparatively (Figure 1), from the present study, *Telfairia pedata* leaf contained higher ash content than *Telfairia Occidentalis* leaf, indicating that the former may likely have more abundant mineral elements than the latter. The result of the ash content of *Telfairia Occidentalis* leaf obtained in this study agreed with the 8.3% ash content of Fluted pumpkin leaf reported [16]. Additionally, the seeds of *Telfairia pedata* have relatively high ash content compared with $2.57 \pm 0.04\%$ obtained by Kim *et al.* [17]. The mean ash content of the two leaves shows a statistically significant dif-

Table 1. Proximate composition of *Telfairia pedata* and *Occidentalis* leaves

Leaves	Moisture (%)	Ash (%)	Fibre (%)	Protein (%)	Lipid (%)	CHO (%)
<i>Telfairia pedata</i>	16.06±0.10	13.30±0.31	19.50±0.031	16.04±0.51	2.00±0.50	49.23±0.79
<i>Telfairia occidentalis</i>	12.20 ±0.20	8.60±0.21	14.68±1.04	22.45±0.5	4.80±0.34	49.44±0.55

ference at ($P < 0.05$). The mean crude protein content of the leaves (Table 3) was significantly different at ($P < 0.05$) and generally higher, probably due to favourable soil fertility and environmental condition. The higher percentage crude protein content of *Telfairia Occidentalis* leaf (22.45 ± 0.50) is in line with (21.14 %) and (22.97 %) reported by Refs. [19] and [14], respectively. Also, the seed of *Telfairia pedata* was found to have contained significantly higher crude protein ($29.08 \pm 1.43\%$) obtained by Kim *et al.* [17] compared with ($16.04 \pm 0.03\%$) obtained in this study. Based on findings by Pearson [20], plants that derive more than 12% of their calorific value from protein are classified as viable protein sources. Therefore, the utilisation of cucurbitaceae leaves in this study suggests that they could be an easily accessible and cost-effective protein source for maintaining regular bodily functions. The result indicates that the *Telfairia Occidentalis* leaf had more protein than the *Telfairia pedata* leaf, probably due to favourable soil fertility and environmental condition.

The lipid content of the leaves was generally low in the studied leaves but higher in the *Telfairia occidentalis* leaf. Adeyeye & Omalojo [16] reported a similar result (3.46%) of the lipid content of the *Telfairia occidentalis* leaf. However, the lipid content (61.20 ± 1.20 %) of the *Telfairia Pedata* seeds obtained by Kim *et al.* [17] is excessively higher than the leaf compared to the present research. The findings suggest that the examined leaves possess health benefits, as their consumption does not appear to contribute to obesity. The mean crude fibre content in (Table 3) of the leaves was significantly different at ($P < 0.05$) and relatively higher. The mean crude fibre of the leaves studied were 19.50 ± 0.51 and 14.51 ± 0.34 for *Telfairia pedata* and *Telfairia occidentalis* leaves, respectively. The crude fibre value of the *Telfairia Occidentalis* leaf agrees with the result (15.40 ± 0.37) obtained [14]. The crude fibre of *Telfairia Pedata* leaf obtained in this research is excessively higher compared with ($1.00 \pm 0.13\%$) obtained by Kim *et al.* [17] in seeds of *Telfairia pedata* (Oyster Nut). This shows that the seeds are generally a poor source of crude fibre. The consumption of dietary fibres has been associated with a reduction in serum cholesterol levels as well as a decreased risk of various health conditions such as coronary heart disease, hypertension, diabetes, colon cancer, and breast cancer [21–23]. From the present analysis, the *Telfairia Pedata* leaf contained a higher percentage of fibre content than the *Telfairia Occidentalis* leaf. However, both the leaves analysed could be a good source of dietary fibre, which could be responsible for the higher carbohydrate content of the leaves.

Carbohydrate content (Table 3) is generally high in all the leaves, and the mean results were not significantly different ($P < 0.05$). The percentage carbohydrate content (49.44 ± 0.55) of *Telfairia Occidentalis* leaf obtained in the present study can be compared favourably with the (51.40 %) carbohydrate contents of *Telfairia Occidentalis* (fluted pumpkin) leaf reported by

Omimakinde [14]. However, the seeds of *Telfairia pedata* (Oyster Nut) contained low carbohydrate content (3.06 %), according to the findings of Kim *et al.* [17]. In retrospect, from this research, the leaf of *Telfairia Pedata* contained an appreciable percentage of carbohydrates (49.23 ± 0.79). The elevated levels of carbohydrates found in the leaves indicate that they contribute substantially to the overall energy content of the food materials. The primary role of carbohydrates is to serve as a source of fuel and energy for the human body, supporting its daily activities and physical exertion.

4. CONCLUSION

In this comparative research, the proximate composition of *Telfairia occidentalis* and *Telfairia pedata* leaves obtained from Katsina metropolis, Nigeria, was carried out, and the results revealed interesting findings. Both leaves portrayed high moisture content, indicating they may not be suitable for prolonged storage because of high water activity. *Telfairia Occidentalis* leaves with higher moisture content will be more susceptible to bacterial spoilage during storage than *Telfairia pedata* leaves. Contrarily, *Telfairia pedata* leaves show higher ash content, indicating a potentially greater abundance of minerals than *Telfairia Occidentalis* leaves. Regarding protein content, *Telfairia Occidentalis* leaves showed significantly higher levels, thus a valuable source of protein suitable for body functions. Although the lipid content of the leaves was relatively low, *Telfairia Occidentalis* leaves contained higher amounts. This suggests that the leaves health-wise are not likely to contribute to obesity when consumed. Furthermore, the crude fibre was significantly higher in *Telfairia Pedata* leaves than in *Telfairia Occidentalis* leaves. The leaves are therefore considered good sources of dietary fibre, which may be linked to their higher carbohydrate content. The carbohydrate content of the leaves was generally high, indicating a significant energy source. The research findings provide valuable insights for understanding these leaves' nutritional value and potential applications, thereby contributing to the broader knowledge of these valuable Cucurbitaceae resources. Further research is recommended to determine the anti-nutrients, minerals and amino acids, and phytochemicals of the leaves for potential nutritive and therapeutic applications.

References

- [1] M. O. Sunmonu, E. O. Ajala, M. M. Odewole, S. Morrison & A. M. Alabi, "Comparative analysis of physico-chemical properties of oil extract from two varieties of fluted pumpkin seeds using different extraction methods", Kathmandu University Journal of Science, Engineering and Technology **13** (2017) 48. <https://doi.org/10.3126/kuset.v13i2.21283>.
- [2] O. R. Temitope, O. O., Olugbenga, A. J. Erasmus, I. Jamilu, M. Y. Shehu, "Comparative study of the physicochemical properties of male and female fluted pumpkin (*Telfairia occidentalis*)", The Journal of Medical Research **6** (2020) 55. https://www.medicinarticle.com/JMR_20202_07.pdf.
- [3] H. D., Irvine, *Commercial Vegetable Growing*, 1st Edt., Oxford University Press, Oxford, UK, 2015.
- [4] A. A. A. Kayode, O. T. Kayode & A. A. Odetola, "*Telfairia occidentalis*

- ameliorates oxidative brain damage in malnourished rats”, *International Journal of Biological Chemistry* **4** (2010) 10. <https://www.cabdirect.org/cabdirect/abstract/20103140081>.
- [5] G. Oboh, E. E. Nwanna & C. A. Elusiyan, “Antioxidant and antimicrobial properties of *Telfairia occidentalis* (Fluted pumpkin) leaf extracts”, *Journal of Pharmacology and Toxicology* **1** (2006) 167. <https://doi.org/10.3923/jpt.2006.167.175>.
- [6] O. A. Eseyin, A. C. Igboasoiki, E. Oforah, P. Ching & B. C. Okoli, “Effects of extracts of *Telfairia occidentalis* leaves on some biochemical parameters in rat”, *Global Journal of Pure Applied Science* **11** (2014) 85. <https://doi.org/10.4314/gjpas.v11i1.16466>.
- [7] A. O. Fasuyi & A. D. Nonyerem, “Biochemical, nutritional and haematological implications of *Telfairia occidentalis* leaf meal as protein supplement in broiler starter diets”, *African Journal of Biotechnology* **6** (2007) 55. <https://www.ajol.info/index.php/ajb/article/view/57100>.
- [8] J. O. Alegbejo, “Production, marketing, nutritional value and uses of fluted pumpkin (*Telfairia occidentalis* Hook. F.) in Africa”, *Journal of Biological Science and Bioconservation* **4** (2012) 20. <https://www.ajol.info/index.php/ajb/article/view/57100>.
- [9] E., Mwakasege, A., Treydte, O., Hoeglenger, N., Kassim, E., Makule, “Variations in nutrient composition of oyster nuts (*Telfairia pedata*) across different agro-climatic conditions”, *Cogent Food & Agriculture* **7** (2021) 1913843. <https://doi.org/10.1080/23311932.2021.1913843>.
- [10] Association of Official Analytical Chemists (AOAC), *Official Methods of Analysis (15th edition)*, Association of Official Analytical Chemists, Wilson Boalevard, Arlington Virginia, USA, 2004, pp 910-928.
- [11] I. T. Gbadamosi, J. O. Moody, A. M. Lawal, “Phytochemical screening and proximate analysis of eight ethnobotanicals used as antimalaria remedies in Ibadan, Nigeria”, *Journal of Applied Bioscience* **44** (2011) 2967. <https://www.m.elewa.org/JABS/2011/44/1.pdf>.
- [12] K. Ihenacho, “Nutritional composition of some leafy vegetables consumed in Imo State, Nigeria”, *Journal of Applied Science and Environmental Management* **13** (2015) 35. <https://doi.org/10.4314/jasem.v13i3.55349>.
- [13] S. S. Nielsen, *Determination of Moisture Content*, In Nielsen, S.S. (Eds.), *Food Analysis Laboratory Manual Food Science Texts Series*, Springer, Boston, USA, 2010. https://doi.org/10.1007/978-1-4419-1463-7_3.
- [14] A. J. Omimakinde, I. Oguntimihin, E. A. Omimakinde, O. Olaniran, “Comparison of the proximate and some selected phytochemicals composition of Fluted Pumpkin (*Telfairia occidentalis*) leaves and pods”, *International Biological and Biomedical Journal* **4** (2018) 206. Accessed on 21/08/2023. https://ibbj.org/browse.php?a_id=196&sid=1&slc_lang=en.
- [15] E. Mwakasege, A. Treydte, O. Hoeglenger, N. Kassim, E. Makule, “Variations in nutrient composition of oyster nuts (*Telfairia pedata*) across different agro-climatic conditions”, *Cogent Food & Agriculture* **7** (2021) 1913843. <https://doi.org/10.1080/23311932.2021.1913843>.
- [16] E. I. Adeyeye & F. O. Omolayo, “Chemical composition and functional properties of leaf protein concentrate of Amaranthus hybrid and *Telfairia occidentalis*”, *Agriculture and Biology Journal of North America* **2** (2011) 499. <https://doi.org/10.5251/abjna.2011.2.3.499.511>.
- [17] M. Y. Kim, E. J. Kim, Y. N. Kim, C. Choi & B. H. Lee, “Comparison of the chemical compositions and nutritive values of various pumpkin (Cucurbitaceae) species and parts”, *Nutrition research and practice* **6** (2012) 21. <https://doi.org/10.4162/nrp.2012.6.1.21>.
- [18] I. C. Onuguh, E. U. Ikhuria & J. U. Obibuzo, “Comparative proximate analysis of *T. occidentalis* (fluted pumpkin) leaves sold in three different markets in Benin City, Nigeria”, *International Journal of Agriculture and Animal Production (IJAAP)* **2** (2022) 7. <https://doi.org/10.55529/ijaap.22.7.12>.
- [19] U. Usonobun & E. Egharebva, “Phytochemical analysis, proximate and mineral composition and in vitro antioxidant activities in *Telfairia occidentalis* aqueous leaf extract”, *Journal of Basic and Applied Science* **1** (2014) 74. Accessed on 21/08/2023. <https://www.biu.edu.ng/blog-post/biujbas-1-1-7>.
- [20] D. Pearson, *Chemical Analysis of Foods*, 7th ed., Churchill, Livingstone, London, 2012, pp 31-105.
- [21] K. Maćkowiak, N. Torlińska-Walkowiak & B. Torlińska, “Dietary fibre as an important constituent of the diet”, *Postepy Hig Med Dosw* **70** (2016) 104. <https://doi.org/10.5604/17322693.1195842>.
- [22] R. U. Ebanu, U. O. Edet, K. I. Anosike, C. A. Etok & T. O. Kanu, “Nutritional analysis and wine production potentials of *Telfairia occidentalis* (fluted pumpkin) leaves and *Cucumis sativus* L.(cucumber) using Baker’s and palm wine yeast strains”, *World News of Natural Sciences* **22** (2019) 12. Accessed on 21/08/2023. <https://agro.icm.edu.pl/agro/element/bwmeta1.element.agro-1a55e233-7840-446b-a678-ee979e9a4144>.
- [23] M. A. China, O. N. Precious & F. Owuno, “Utilisation of fluted pumpkin (*Telfairia occidentalis*) seed milk for the production of textured vegetable protein”, *European Journal of Agriculture and Food Sciences* **3** (2021) 4. <https://doi.org/10.24018/ejfood.2021.3.4.81>.